

## Why the product rule is true:

$$\begin{aligned} & \frac{d}{dx} (f(x) g(x)) \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) g(x+h) - f(x) g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) g(x+h) - \color{red}{f(x) g(x+h)} + \color{red}{f(x) g(x+h)} - f(x) g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( \frac{\color{blue}{f(x+h) g(x+h)} - \color{blue}{f(x) g(x+h)}}{h} + \frac{\color{blue}{f(x) g(x+h)} - \color{blue}{f(x) g(x)}}{h} \right) \\ &= \lim_{h \rightarrow 0} \left( \left[ \frac{\color{blue}{f(x+h)} - \color{blue}{f(x)}}{h} \right] \color{blue}{g(x+h)} + \color{blue}{f(x)} \left[ \frac{\color{blue}{g(x+h)} - \color{blue}{g(x)}}{h} \right] \right) \\ &= \color{blue}{f'(x)g(x)} + \color{blue}{f(x)g'(x)} \end{aligned}$$

## Why the quotient rule is true:

Let  $h(x) = \frac{f(x)}{g(x)}$ . We want to find  $h'(x)$

$$f(x) = g(x)h(x)$$

$$\Rightarrow f' = g'h + gh' \quad \text{by the Product Rule}$$

$$f' = g' \left( \frac{f}{g} \right) + gh'$$

$$f'g = g'f + g^2h'$$

$$f'g - g'f = g^2h'$$

$$h' = \frac{f'g - g'f}{g^2}$$

1. Find the derivative of each function

(a)  $h(x) = x^{32} \ln(x)$

(c)  $h(x) = \frac{3 + 2x^{-3}}{8x^3 - 4x}$

(b)  $h(x) = \sqrt{x} \left( \ln(x) + \frac{2}{x^3} \right)$

(d)  $h(x) = x \ln(x) - x$

2. Find an antiderivative for each function

(a)  $f(x) = 2x \sin(x) + x^2 \cos(x)$

(b)  $f(x) = \ln(x) + \frac{1}{x}$

3. Find the derivative of each function and verify your answer by graphing

(a)  $g(x) = \sin(x^2)$

(b)  $g(x) = (\sin(x))^2$