Why the product rule is true:

$$\frac{d}{dx}\left(f(x)\,g(x)\right)$$

$$= \lim_{h\to 0} \frac{f(x+h)\,g(x+h) - f(x)\,g(x)}{h}$$

$$= \lim_{h\to 0} \frac{f(x+h)\,g(x+h) - f(x)\,g(x+h) + f(x)\,g(x+h) - f(x)\,g(x)}{h}$$

$$= \lim_{h\to 0} \left(\frac{f(x+h)\,g(x+h) - f(x)\,g(x+h)}{h} + \frac{f(x)\,g(x+h) - f(x)\,g(x)}{h} \right)$$

$$= \lim_{h\to 0} \left(\left[\frac{f(x+h) - f(x)}{h} \right] g(x+h) \;+\; f(x) \left[\frac{g(x+h) - g(x)}{h} \right] \right)$$

$$= f'(x)g(x) + f(x)g'(x)$$

Why the quotient rule is true:

Let $h(x) = \dfrac{f(x)}{g(x)}$. We want to find $h'(x)$

$$f(x) = g(x)h(x)$$

$$\Rightarrow \quad f' = g'h + gh' \qquad \text{by the Product Rule}$$

$$f' = g'\left(\frac{f}{g}\right) + gh'$$

$$f'g = g'f + g^2h'$$

$$f'g - g'f = g^2h'$$

$$h' = \frac{f'g - g'f}{g^2}$$

Work on these with your partner(s) at the board

1. Find the derivative of each function

   (a) $h(x) = x^{32} \ln(x)$

   (b) $h(x) = \sqrt{x}\left(\ln(x) + \dfrac{2}{x^3}\right)$

   (c) $h(x) = \dfrac{3 + 2x^{-3}}{8x^3 - 4x}$

   (d) $h(x) = x\ln(x) - x$

2. Find an antiderivative for each function

   (a) $f(x) = 2x\sin(x) + x^2\cos(x)$ (b) $f(x) = \ln(x)$

3. Find the derivative of each function and verify your answer by graphing

   (a) $g(x) = \sin(x^2)$ (b) $g(x) = (\sin(x))^2$